
 <div style="text-align: center;"> Oregon Health & Science University Hospital and Clinics Provider's Orders </div> <div style="text-align: center; margin-top: 10px;">  </div> <div style="text-align: center; margin-top: 10px;"> <small>ADULT AMBULATORY INFUSION ORDER</small> Alteplase (t-PA) Infusion for Dialysis Catheters </div> <div style="text-align: center; margin-top: 20px;"> <small>Page 1 of 2</small> </div>	<div style="margin-bottom: 10px;">ACCOUNT NO. _____</div> <div style="margin-bottom: 10px;">MED. REC. NO. _____</div> <div style="margin-bottom: 10px;">NAME _____</div> <div style="margin-bottom: 10px;">BIRTHDATE _____</div> <div style="text-align: right; margin-top: 20px; font-style: italic;">Patient Identification</div>
ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.	

Weight: _____ kg Height: _____ cm

Allergies: _____

Diagnosis Code: _____

Treatment Start Date: _____ Patient to follow up with provider on date: _____

****This plan will expire after 365 days at which time a new order will need to be placed****

NURSING ORDERS:

1. Aspirate 3 mL of blood from each dialysis lumen to remove high dose heparin prior to flushing
2. Refer to nursing and IV therapy guidelines for care of central venous catheters
3. Follow facility policies and/or protocols for vascular access maintenance with appropriate flush solution, dec clotting (alteplase), and/or dressing changes.

MEDICATIONS:

INFUSION ORDERS

LUMEN #1

- ☐ alteplase (ACTIVASE) 2 mg in sodium chloride 0.9% 100 mL, intracatheter, ONCE over 4 hours as needed for occluded dialysis catheter lumen (Maximum of 4 mg total in all lumens)

LUMEN #2

- ☐ alteplase (ACTIVASE) 2 mg in sodium chloride 0.9% 100 mL, intracatheter, ONCE over 4 hours as needed for occluded dialysis catheter lumen (Maximum of 4 mg total in all lumens)

POST INFUSION ORDERS

LUMEN #1

- ☐ alteplase (ACTIVASE) 2 mg, intracatheter, ONCE, Label dressing "TPA dwell" with date, time, and RN initials

OR

- ☐ heparin 1000 units/mL, 1-5 mL, intracatheter, ONCE, Pack dialysis catheter with the volume of catheter plus 0.25 mL

LUMEN #2

- ☐ alteplase (ACTIVASE) 2 mg, intracatheter, ONCE, Label dressing "TPA dwell" with date, time, and RN initials

OR

- ☐ heparin 1000 units/mL, 1-5 mL, intracatheter, ONCE, Pack dialysis catheter with the volume of catheter plus 0.25 mL



Oregon Health & Science University
Hospital and Clinics Provider's Orders

ADULT AMBULATORY INFUSION ORDER
**Alteplase (t-PA) Infusion for Dialysis
Catheters**

Page 2 of 2

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.

HYPERSENSITIVITY MEDICATIONS:

1. NURSING COMMUNICATION – If hypersensitivity or infusion reactions develop, temporarily hold the infusion and notify provider immediately. Administer emergency medications per the Treatment Algorithm for Acute Infusion Reaction (OHSU HC-PAT-133-GUD, HMC C-132). Refer to algorithm for symptom monitoring and continuously assess as grade of severity may progress.
2. diphenhydramine (BENADRYL) injection, 25-50 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
3. EPINEPHrine HCl (ADRENALIN) injection, 0.3 mg, intramuscular, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
4. hydrocortisone sodium succinate (SOLU-CORTEF) injection, 100 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
5. famotidine (PEPCID) injection, 20 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction

By signing below, I represent the following:

I am responsible for the care of the patient (*who is identified at the top of this form*);

I hold an active, unrestricted license to practice medicine in: ☐ Oregon ☐ _____ (*check box that corresponds with state where you provide care to patient and where you are currently licensed. Specify state if not Oregon*);

My physician license Number is # _____ (MUST BE COMPLETED TO BE A VALID PRESCRIPTION); and I am acting within my scope of practice and authorized by law to order Infusion of the medication described above for the patient identified on this form.

Provider signature: _____ **Date/Time:** _____

Printed Name: _____ **Phone:** _____ **Fax:** _____

INFUSION REFERRAL TEAM Phone (providers only) (971) 262-9645 Fax completed orders to (503) 346-8058 Infusion orders located at: www.ohsuknight.com/infusionorders	<input checked="" type="checkbox"/> Please indicate the patient's preferred clinic location below	
	<input type="checkbox"/> BEAVERTON OHSU Knight Cancer Institute	15700 SW Greystone Court Beaverton OR 97006
	<input type="checkbox"/> NW PORTLAND Legacy Good Samaritan campus	Medical Office Building 3 – Suite 150 1130 NW 22nd Ave, Portland OR 97210
	<input type="checkbox"/> GRESHAM Legacy Mount Hood campus	Medical Office Building 3 – Suite 140 24988 SE Stark, Gresham OR 97030
	<input type="checkbox"/> TUALATIN Legacy Meridian Park campus	Medical Office Building 2 – Suite 140 19260 SW 65th Ave, Tualatin OR 97062